

Accepted Manuscript

Organic Reactions for the Electrochemical and Photochemical Production of CO₂ to Fuels-The Reduction Chemistry of Carboxylic Acids as Bent CO₂ Surrogates, Aldehydes, Alcohols and Alkyls

Oana R. Luca, Aidan Q. Fenwick

PII: S1011-1344(15)00138-4

DOI: <http://dx.doi.org/10.1016/j.jphotobiol.2015.04.015>


Reference: JPB 10012

To appear in: *Journal of Photochemistry and Photobiology B: Biology*

Received Date: 17 January 2015

Revised Date: 10 April 2015

Accepted Date: 16 April 2015



Please cite this article as: O.R. Luca, A.Q. Fenwick, Organic Reactions for the Electrochemical and Photochemical Production of CO₂ to Fuels-The Reduction Chemistry of Carboxylic Acids as Bent CO₂ Surrogates, Aldehydes, Alcohols and Alkyls, *Journal of Photochemistry and Photobiology B: Biology* (2015), doi: <http://dx.doi.org/10.1016/j.jphotobiol.2015.04.015>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Organic Reactions for the Electrochemical and Photochemical Production of CO₂ to Fuels-
The Reduction Chemistry of Carboxylic Acids as Bent CO₂ Surrogates, Aldehydes,
Alcohols and Alkyls**

Oana R. Luca,^{a§} Aidan Q. Fenwick^a

[§]*corresponding author*

^a*Joint Center for Artificial Photosynthesis, Division of Chemistry and Chemical Engineering,
California Institute of Technology, Pasadena, California 91125, United States.*

Abstract

The present review highlights the organic chemistry involved in the reduction of CO₂ and related intermediates to chemical fuels. We focus on known chemical reactions of CO₂ with organic molecules to yield carboxylic acid derivatives that create opportunities for reductive tandem electrochemical/chemical processes emerge. We draw parallels between long-standing knowledge of CO₂ reactivity from organic chemistry, organocatalysis, surface science, electrocatalysis and theoretical calculations related to these processes. We also point out some possible non-faradaic chemical reactions that could contribute to observed product distributions in the production of solar fuels from CO₂.

1. Introduction

2. CO₂ Reduction Chemistry and Intermediates in Solar Fuel Production

2.1. CO₂ Reduction at Cu – Intermediates

2.2. Catalytic Cycle with Molecular Preactivation – Intermediates and Functional-Group Interconversion Overview

3. Activation Chemistry of CO₂: Formation of Carboxylic Acids

3.1. Photosynthesis: RuBisCo and the Calvin Cycle

3.2. C1-Carriers in Nature: Folate Chemistry

3.3. CO₂ Binding to Organic Bases in the Absence of an Added Lewis Acid

3.3.1. N-Heterocyclic Carbene (NHC) Carboxylates and Catalysis

3.3.2. N-Heterocyclic Olefin (NHO) Carboxylates and Catalysis

3.3.3. Carbamates and Related Reductions

3.3.4. Bicarbonates and Related Reductions

3.4. Lewis-Acid/Base Activation and Catalysis

3.5. CO₂ Binding to Metal Centers

3.5.1. Binding Modes

3.5.2. Metal *vs* Ligand Coordination and Redox-Active Ligands

3.5.3. Electrocatalytic CO₂ Reduction *via* Metal Carboxylates

3.6. Electrocarboxylation Reactions

4.0. Reduction of Carboxylic Acids to Aldehydes

4.1. Carboxylic Acid Reductase

4.2. Chemical - Carbamate reduction

4.3. Electrochemical Reduction

5.0. Reduction of Carboxylic Acids and Carbamates to Alcohols

5.1. LiAlH_4 , NaBH_4 and BH_3

5.3. Bouveault-Blanc Reductions

5.4. Electrochemical Bouveault-Blanc Reductions

5.5. Reduction of NHC Carboxylates with Silanes – Methanol production

5.6. Other Electrochemical Reductions

6.0. Reduction of Carboxylic Acids to Alkyls

6.1. Carboxylic Acid Reductase Tandem catalysis

6.2. Barton-McCombie Decarboxylation

6.3. Electrochemical Reduction

7.0. Reduction of Aldehydes to Alcohols

7.1. NaBH_4 and LiAlH_4

7.2. Crossed Cannizzaro reaction: Formaldehyde as a reducing agent

7.3. Silanes, Boranes and NHC Boranes

7.4. Electrochemical Reduction: Reductive Dimerization.

7.5. Electrochemical Reduction to Primary Alcohols without Dimerization on Cu Electrodes

8.0. Reduction of Aldehydes to Alkyls

8.1. Wolff-Kishner Reduction

8.2. Caglioti Reaction: Reduction of Tosylhydrazones

8.3. Clemmensen Reduction

8.4. Eschweiler Clark Reduction

9.0. The Reduction of Alcohols to Alkyls

9.1. Barton-McCombie Deoxygenation

9.2. Electrochemical Deoxygenation

10.0 Conclusions

1. Introduction

Carbon dioxide (CO_2) is a linear molecule that has attracted attention in recent years as a source of renewable energy through its conversion to chemical fuels[1]. The current discussion will highlight the use of molecular preactivators to alter the structure of the CO_2 molecule for selective access to high-value fuels *via* chemical and electrochemical means. We emphasize the chemical transformations of directly relevant functional groups and point out chemical reactions related to the observed chemistry.

In order to achieve the efficient conversion of CO_2 to chemical fuels, a catalyst is generally required to lower the activation energy of the chemical reduction, and many reaction pathways with comparable activation energies or shared reaction intermediates are generally accessible, leading to a variety of reaction products during catalysis. Although significant effort has been specifically centered on catalyst development for reducing CO_2 in solution, this overview aims to draw attention to examples of CO_2 -bending reactions with organic bases to form carboxylate intermediates as starting points for reactions related to CO_2 conversions. Under this paradigm, the reduction of CO_2 becomes an exercise in the electrochemical reduction of carboxylic acids and derivatives, where a wealth of knowledge from organic chemistry comes into play.

The use of substoichiometric concentrations of certain organic molecules and their intrinsic reactivity as catalysts or mediators for chemical processes defines the field of organocatalysis [2, 3]. At the same time, these transformations are directly reminiscent of reactions in nature, where cofactors and coenzymes mediate biological pathways[4]. Such organic-mediated transformations with regards to the conversion of CO_2 to chemical fuels have been reviewed by Hu and coworkers[5]. The work summarizes the use of tetraalkylammonium salts[6], ionic liquids[7], and pyridinium derivatives[8] that have been successfully shown to act as mediators in the selective (photo-) electrochemical reduction of CO_2 to products such as CO and methanol.




Figure 1. Opportunities for Tandem Chemical/ Electrochemical CO₂ reduction to fuels. [1]

We recognize that the organic reductive chemistry of carbonyl-containing moieties generally involves the formation of an activated adduct as a first step. Transformations of this kind, such as the Wolff-Kishner (Sections 8.1) and the related Caglioti reaction (Section 8.2) and as two examples, involve the formation of an activated intermediate poised to undergo further reduction.

This particular strategy is an opportunity for new directions in the electrochemical reduction of CO₂ (Figure 1), where chemical transformations to form more easily reduced species could be performed in tandem with electrochemical transformations, when needed. Given this approach, to the best of our knowledge, reactions such as the Eschweiler-Clark (Section 8.5) and Crossed Cannizzaro (Section 7.2) have not ever been addressed as possible contributors to observed product distributions in current efforts in electrocatalysis for CO₂ conversion to fuels.


We limit the current discussion to organic-mediated, stoichiometric transformations and their mechanisms. Several of these reactions have been performed through transition-metal catalyzed hydrogenation and transfer hydrogenations. These reactivities remain outside the scope of the current discussion as they have been described elsewhere.[9-13]

2. CO₂ Reduction Chemistry and Intermediates in Solar Fuel Production

Organic molecules regenerated at electrode surfaces have been shown to perform energy-relevant redox chemical transformations [14] [3] [5]. Such molecules have also been involved in

metal/ligand cooperativity in electrocatalytic dehydrogenation reactions with molecular precursors. No activity was observed in the absence of the redox-active component[15, 16].

Scheme 1 depicts a reduction pathway from a carboxylic acid down to an overall 8 H^+ and 8 e^- reduced species: CH_4 . This particular scheme highlights opportunities for tandem conversion processes, catalytic or otherwise.




Scheme 1. General CO_2 reduction pathway *via* a Preactivation motif: *i* Prebending of the CO_2 prior to any chemical transformation.

During the preactivation step (*i*) the CO_2 molecule is converted to a carboxylic acid. A carboxylic acid (RCOOH) can then be converted to an aldehyde (RCHO) by 2 H^+ and 2 e^- (*ii*) which can subsequently be reduced to an alcohol (RCH_2OH) (*iii*) and then further to an alkyl (RCH_3) (*iv*) which could release the carbon fragment as CH_4 in a R-C bond-breaking event (*v*).

2.1. Intermediates in the CO_2 Electroreduction at Cu Surfaces

Heterogeneous electrocatalysis of the CO_2 reduction reaction comprises one of the largest technology gaps for efficient, sustainable CO_2 reduction and as such, the work discussed herein is a novel direction for the field and will likely enable the deployment of efficient, sustainable fuel generation. Homogeneous electrocatalysis has the advantage of tunability of the catalyst at the molecular level, however the operating conditions render current state-of-the-art systems technologically irrelevant[17]. Coupling the electrocatalytic reactions (both homogeneous and heterogeneous) with sorption and preconversion of CO_2 from the atmosphere will likely yield discoveries that translate from basic materials discovery, through cathode/catholyte components, to design of test-bed prototypes in the future.

In computational work by Nørskov [18] and coworkers, sequential reduction steps from CO_2 to methane are addressed as depicted in Scheme 2. Comparisons between enzymatic CO_2 reduction activities with heterogeneous metal surface [19] electrocatalysis bring forward a rate-determining step involving a CO-bound intermediate.



Scheme 2. Electrochemical Reduction of CO₂ to methane –theoretical calculations[18].

The mechanistic work proposes a series of chemical transformations as depicted in Scheme 1 and 2 with the formation of a carboxylate intermediate (**1**) as the first step in the process. In this particular regime, the carboxylate can then undergo a dehydration reaction to yield a bound [Cu-CO] (**2**) fragment, which then can be further reduced to a formyl fragment [Cu-CHO] (**3**) followed by an additional reduction to an O-bound formaldehyde species [Cu-OCH₂] (**4**) and then to a methanolic unit [Cu-OCH₃] (**5**) that can release CH₄ upon further reduction[18].

2.2. Catalytic Cycle with Molecular Preactivation – Intermediates and Functional-Group Interconversion Overview

Given the context of heterogeneous electrocatalysis in Section 2.1, our current discussion will focus on reductive functional group interconversions between carboxylic acids, aldehydes, alcohols and alkyls and we highlight examples of reactions involving both chemical and electrochemical regimes as can be seen in Figure 2. The related conversions and sections of this review are summarized in Table 1.





Figure 2. Chemical vs electrochemical conversions laid out in this review.

Table 1. Overview of reactivity discussed in the present work.

<div>Products</div> <div>Starting Materials</div>	RCOOH	RCHO	RCH₂OH	RCH₃	RH
RCOOH		Section 4	Section 5	Section 6	Section 6
RCHO			Section 7	Section 8	n/a
RCH₂OH				Section 9	n/a
RCH₃					n/a
RH					

3. Activation Chemistry of CO₂: Formation of Carboxylic Acids

3.1. Photosynthesis: RuBisCo and the Calvin Cycle




Scheme 3. CO₂ activation by ribulose 1,5-phosphate as the first step in Ribulose-1,5-bisphosphate carboxylase (RuBisCo) carbon fixation.

In process of natural photosynthesis, photosystem I is the membrane-bound unit where CO₂ is captured from the atmosphere and converted to sugars. The Calvin cycle is the metabolic pathway of CO₂ capture and conversion of sugars that takes place in the stroma of the chloroplast. RuBisCo – (Ribulose-1,5-BISphosphate Carboxylase/Oxygenase) is the enzyme responsible for the carbon fixation mechanism as described in Scheme 3. RuBisCo is believed to be the most abundant protein on Earth [20, 21].

The process occurs through a series of steps using ATP as an energy source and NADPH₂ as terminal reductant. The first step is the step of interest in the present discussion: carbon capture. The second and third phases of the cycle use the reducing equivalents from NADPH₂ and ATP to form sugars (step 2) and then regenerate RuBisCo in step 3. The net stoichiometry of the overall Calvin cycle suggests that for every three molecules of fixated CO₂, one molecule of 3-phosphoglycerate (3PG, 9) is formed at the expenditure of nine molecules of ATP and six NaDPH₂.

As can be seen in the first step, the CO₂-trapping event occurs *via* a nucleophilic attack of an enediolate (7) onto CO₂ to yield a β-ketoacid (8). After a sequence of steps, the bound CO₂ gets converted to energy-rich molecules such as glucose.

3.2. C1-carriers in Nature: Folate Chemistry




Scheme 4. Hydrofolate intermediates as C1 carriers in nature [22, 23].

Folates are N-containing chemical compounds responsible for aiding in the interconversion of formyl fragments to methyl in biological systems. They act as C1 carriers between several oxidation states of the carbon-containing organic fragment. The observed structures (Scheme 4) show several intermediates in folate metabolism including a range of N-bound methyl, formyl, methylene, methenyl and formimino fragments.

Such chemical transformations have been highlighted in reports related to tetrahydrofolate and tetrahydromethanopterin as different C1 carriers[22, 24]. The role of such conversions have been suggested in methanogenesis pathways[25-27]. Interestingly, a report on the involvement of such intermediates in the conversion of acetate into methane has also been made available[28].

3.3. CO₂ binding to Organic Bases in the Absence of an Added Stabilizing Lewis Acid

3.3.1. N-Heterocyclic carbene (NHC) Carboxylates



N-heterocyclic carbenes (NHC, **10**) have had a long-standing role in transition metal catalysis as versatile ligand frameworks with easily modulated donating properties[29]. The advent of organocatalysis has now exploited the organic fragments themselves as catalysts for a myriad of organic transformations in metal-free conversions[30, 31]. Most excitingly, the N-heterocyclic carbene fragments have been shown to react with CO₂ to form zwitterionic imidazolium carboxylates (Eq 1, **11**)[32]. These compounds were first reported as a protection method for air-sensitive NHC fragments and as precursors for NHC ligands for the synthesis of transition-metal complexes.

Louie and coworkers have now shown that the strength of binding CO₂ to the NHC fragment is tunable in a series of decarboxylation and transcarboxylation studies[33, 34]. Thermal stability studies have yielded a series of data shown in Figure 3[35, 36]. From the available data, we have observed a trend in the temperature (T) of CO₂ release of these zwitterionic species. The T correlates linearly with the dihedral angle between the CO₂ fragment (red) and the NHC fragment (blue) of these zwitterionic species. The zwitterions with the more electron-donating and less sterically-hindered NHCs **12** and **13** are almost flat, thus suggesting strong binding and possible conjugation with partial double-bond character between the CO₂ fragment and the NHC.





Figure 3. Tuning of the strength of CO₂ binding with N-heterocyclic carbenes. Plots of the T of CO₂ release vs dihedral angle between NHC and CO₂[35].

Although outside of the scope of the current discussion as it relates to an oxidative process, we would like to note an example of electrode-driven organocatalytic process[37]. Bodyston and coworkers report the preactivation of an aldehyde *via* a NHC organic fragment. The preactivation is then followed by an anodic process to electrochemically generate an oxidized intermediate than can react with an alcohol to form an ester.


► In the reductive regime, the organocatalytic reduction of CO₂ to methanol has now been reported by Riduan *et al* (Eq 2)[38]. The reaction is mediated by a free NHC binding the CO₂ and initial formation of an NHC-CO₂ adduct (**11**). The adduct has been shown to be able to undergo sequential reductions by the terminal reducing agent, the silane. Intermediate formates (**20**) and silanols (**21**) have been observed by mass spectrometry.



Scheme 5. Silane as a terminal reducing agent in NHC-catalyzed CO_2 reduction to methanol.


Upon aqueous workup, the terminal silanol reduced species is able to release methanol as a $6e^-$ reduced *liquid* product. Mechanistic studies [39] [40] suggest that the limiting step in the chemistry is the formation of strong Si-O bonds. This provided a chemical opportunity for replacing the silane reducing agent with an electrode-generated reducing agent, shown to take the reduction of the NHC-carboxylate down to an $8e^-$ reduced product, methane, as reported by Luca *et al*[41]. To the best of our knowledge, this is the first report of such a reductive transformation and we remain forward-looking to developments in the field.

3.3.2. N-Heterocyclic Olefin (NHO) Carboxylates and Catalysis



N-heterocyclic olefins (**22**) are a class of Lewis-basic species that have been highlighted for their ability to stabilize main group hydrides in usually low oxidation states[42]. In addition, borenium cation stabilization has also been shown[43]. Showing a similar behavior with the NHCs, they have been shown to be sufficiently Lewis-basic to react and bend CO_2 (Eq 3) to form adducts such as **23**.

NHOs have been shown to catalyze polymerization reactions[44]. Similar to the NHC chemistry in 3.3.1., NHO carboxylates have formed through productive CO_2 -binding and have now emerged as competent catalysts for CO_2 transfer (Scheme 6). They were shown to react with propargylic alcohols (**24**) to form α -alkylidene cyclic carbonates in high yields (**25**)[45].



Scheme 6. N-heterocyclic olefin-catalyzed carboxylation of alkynes.


The authors note that the activity of NHO- CO_2 adducts is in some cases 200-fold when compared to the NHC- CO_2 analogues (Section 3.3.1). The hypothesis for the particular trend was the ability of the NHO- CO_2 to release the CO_2 fragment easily due to its less Lewis basic nature. This particular property may become an important factor in the development of electrode-driven processes.

3.3.3. Carbamates and Related Reductions (1)

Carbon dioxide has been shown to react with amines [46-48] (Eq 4, **26**) and related derivatives to form carbamates (**27**). The chemical reduction of carbamate-related species has

been recently reviewed elsewhere[49]. Highly nucleophilic primary amines have been shown to react with CO₂ to form urea derivatives in a formal 2e⁻ reduction process[50].

The less-nucleophilic pyridines have also been shown to react with CO₂ anion radicals[51] to form carbamate anion radicals. This chemistry has been the protagonist of a series of electrochemical and photoelectrochemical studies for the pyridine-mediated conversion of CO₂ to chemical fuels[8, 52].




An important report in the field of CO₂ preactivation comes from Dyer and coworkers. [53] 6,7-dimethyl-4-hydroxy-2-mercaptopteridine, a N-containing commercially-available heterocyclic compound was shown to act as a mediator for the electrochemical reduction of CO₂ to methanol, albeit with modest faradaic efficiencies (10-23%). Mass spectrometry indicates that the chemistry may be facilitated by a carbamate formed *in situ*. Other observed products during the catalysis are formates and aqueous formaldehyde. Reduction of 1-pyrrolyl and 1-indolyl carbamates to hemiaminals[54] has also been demonstrated electrochemically.

We believe that side chemical reactions between products of CO₂ reduction, particularly formaldehyde and formic acid under the conditions of electrocatalytic conversion may have important contributions to the products observed. For example, in an attempt to perform the electrochemical conversion of CO₂ bound to an amine preactivator under basic aqueous conditions, an Eschweiler- Clark [55] *chemical* reaction is favored to occur. This reaction may be responsible for observed product distributions. We highlight this type of transformation in Section 8.5.

3.3.4. Bicarbonates and Related Reductions

Hydroxide anions react with carbon dioxide in a similar fashion to amines and N-heterocyclic carbenes to form bicarbonate anions (Eq 5, **28**). This type of reactivity is of particular interest in the field of solar fuels, as several of the proposed solar-fuel devices involve aqueous conditions in basic media due to limitations on the anodic side chemistry of a solar-fuel device[1].




Bicarbonate reduction is an important discussion topic, as current solar-fuel device designs will likely rely on basic conditions. Under such conditions, the CO₂ concentration in an aqueous phase is increased, as CO₂ speciates in solution as bicarbonate [1]. Most excitingly a reversible bicarbonate/formate interconversion has been the protagonist of a recent report illustrating the

concept of a formate battery[56]. This is an example of a Virtual Hydrogen Storage device [14, 57, 58] using CO₂ as a molecular framework for the storage of hydrogen equivalents as highlighted in a recent report[1].

3.4. Lewis-Acid/Base Activation and Catalysis


A Lewis acid, such as a borane, can poise the CO₂ molecule to more easily accept reducing equivalents, such as hydrides. (30). A Lewis base (29) is then able to aid in the formation of a CO₂ complex in the form of a LB⁺CO₂LA⁻ (31)[59]. In Scheme 7, such binding is demonstrated to occur with a phosphine Lewis Base and a borane Lewis Acid. Catalytic reductions of CO₂ to methanolic species have been shown with the help of phosphine-borane (33) binding and catalysis [60, 61] with a borane terminal reducing agent (32).



Scheme 7. Lewis acid-base activation and catalyzed reduction of CO₂ with phosphine boranes[60, 61].

In a similar fashion, aided by Frustrated Lewis-Acid/Base Pair (FLP), the selectivity for CO₂ reduction was shown to switch to a fully reduced 8e⁻ product, methane, in the case of an amine-borane hydrosilylation [62] and aluminum-silane CO₂ reduction catalysis [63]. An important application of these FLPs is the activation of H₂ and subsequent use in catalytic hydrogenation reactions, [64] some of which involve carbonyl groups [65].

To illustrate a relevant chemical transformation, we describe a silane-borane FLP motif and subsequent CO₂ binding (Scheme 8, 34). The initial binding and activation of CO₂ leads to the formation of a formate species (35). This event is followed by a cascade hydrosilylation reduction [60]. The frustrated Lewis pair between a borane and a silane has now been directly observed [66] and several instances of related asymmetric hydrogenation and hydrosilylation have now been reviewed[67].



Scheme 8. Lewis acid-base activation and catalyzed reduction of CO₂ with silane boranes.

3.5. CO₂ Binding to Metal Centers

3.5.1. Binding Modes

Next in our discussion, we illustrate some binding modes observed in organometallic and coordination compounds and relate these modes with the binding of organic molecules to CO₂. An extensive discussion of such binding has been made available by Gibson [68] and Leitner [69] and we illustrate some structural examples in Figure 4.





Figure 4. Binding Modes of CO₂.

There are five observed binding modes of CO₂ to metal centers (Figure 4). The coordination type depends on the nature of the metal, on the oxidation states and, consequently, on the available orbitals of the metal center. The explicit coordination environment of the metal complex also plays an important role. If the metal complex has available empty orbitals, coordination with donation of electron density from the O ligand will be possible (end-on for **37**, **39** and **40** and side-on for **38** in Figure 4). CO₂ binding to Lewis acidic metals usually involves coordination through the oxygens and depending on the available anchoring sites and oxidation states, it can occur η¹-CO₂ (**39**) or η²-CO₂ (**40**).

If the metal complex has a low-valent nucleophilic site, then CO₂ coordination occurs as an η²-CO₂ complex with the metal bound to the central carbon to form a carboxylate (**36**). This particular type of coordination is also observed in the presence of organic nucleophiles as

described in Section 3.3 and is of particular interest in the field of electrocatalysis for CO₂ reduction with molecular species (section 3.4.3). The central C-binding mode (**36** in Figure 4) of CO₂ is the protagonist of several of the state-of-the art electrocatalytic processes for CO₂ conversions at metal centers[1, 70]. An example of such reactivity is given in Section 3.5.3.

3.5.2. Metal vs Ligand Coordination and Redox-Active Ligands[71]




A topic of great interest in the catalysis community, the concept of redox-active actor ligands, plays an important role in CO₂ coordination at metal centers and has extensively been described elsewhere[71]. We give one example of ligand participation in CO₂ binding to the organic fragment of a Ru complex with a redox-active ligand (Eq 6).

Milstein and coworkers have now shown that the presence of a nucleophilic redox-active diphosphine ligand on a Ru center (**41**) facilitates the indirect reduction of carbonates under hydrogenation conditions[72]. Specifically, the first step in this chemistry is the coordination of CO₂ to the relevant metal complex yields an intermediate with the CO₂ fragment C-bound to the nucleophilic ligand center and O-bound to the Ru in a side-on fashion (**42**).

3.5.3. Electrocatalytic CO₂ reduction *via* Metal Carboxylates

As CO₂ electrocatalysis [1] for the formation of fuels remains a topic of great interest, we illustrate one of many examples of CO₂ binding to a reduced metal complex; specifically, we make note of an important intermediate in the Re-catalyzed conversion of carbon dioxide to CO.




Scheme 9. Electrocatalytic CO₂ reduction cycle with Re(bpy)(CO)₃X.[70]

In the current catalytic understanding, binding mode **36** occurs after the reaction of CO₂ with a nucleophilic reduced complex **43**. This carboxylate **44** can then be protonated and further reduced in a catalytic cycle as shown in Scheme 9. A Mn variant of this chemistry has also been reported [73] and specifically, the reduction of a (bpy)(CO)₃Mn^ICO₂H was observed *via* IR-spectroelectrochemistry[74].

3.6. Electrocatalytic Reactions


An electrocarboxylation reaction is the reaction of CO₂ with an organic molecule to form carboxylic acids under electrochemical conditions[75]. The preparative and large-scale implementation of such reactions has been extensively studied and the results have been described in detail elsewhere[76].



Scheme 10. Electrocatalytic cycle for the electrocarboxylation of alkynes via $\text{Ni}(\text{bpy})_2^0$ in a single-compartment electrolysis cell. [75]

To describe this type of electrode-induced CO_2 preactivation, we give an example of electrochemical incorporation of CO_2 into unactivated alkynes (**46**). This reaction has been successfully mediated by an electro-generated $\text{Ni}(\text{bpy})_2^0$ (**45**) in a single-compartment electrolysis cell to yield α,β -unsaturated carboxylic acids (**48**, Scheme 10). In the case of 4-octyne, a nickelacycle (**47**) was isolated and characterized[75].

A further example of CO -selective CO_2 -reduction molecular electrocatalyst is $\text{Ni}(\text{cyclam})^{2+}$ [77, 78]. Interestingly, $\text{Ni}(\text{cyclam})^{2+}$ has also been shown to be an effective mediator for the electrosynthesis of cyclic carbonates (**50**) from epoxides (**49**) and CO_2 (Eq 7).




4.0. Reduction of Carboxylic and Carbamic Acids to Aldehydes

4.1. Carboxylic Acid Reductase

The reduction of carboxylic acids to aldehydes can be performed enzymatically by carboxylic acid reductase (CAR) [79, 80]. The enzyme is able to convert non-activated carboxylates to the corresponding reduced aldehydes without overreduction to the corresponding alcohols [79, 80].

Specifically, studies on recombinant CAR have demonstrated conversion of fatty acids (C_6 - C_{18}) as well as benzoic, vanillic, and ferulic acids to their corresponding aldehydes. The broad substrate specificity of CAR, in combination with an aldehyde reductase co-enzyme, has been studied as possible metabolic route for the biosynthesis of fuels and commodity chemicals. Since the work is biochemical in nature, it falls outside of the scope of this discussion, although it sets an important precedent for the chemistry discussed in Section 9.

4.2. Chemical – Carbamate Reductions



The preactivation of CO_2 with amines was described in Section 3.5.3. In a recent study, 1,5,7-Triazabicyclo[4.4.0]dec-5-ene was shown to catalyze amine hydroformylation (Eq 8). The CO_2 fragment is first stabilized and activated into a zwitterionic carbamate (**51**). This species is able to undergo reductive hydrosilylation to the corresponding aldehyde species (**52**).

An in-depth study on this type of organocatalytic formylation of morpholine using CO_2 and silanes is now in the literature [81] [60]. This chemistry comes as an alternative to previous methods employing Ru catalysts [82, 83]. If the terminal silane reducing agent would be electrochemically


4.3. Electrochemical

Although the reduction of a free carboxylic acid directly to an aldehyde product has not been reported to the best of our knowledge, we point out the conversion of carboxylic acid derivatives such as esters to the corresponding aldehydes under electrochemical conditions. Alkylbenzoate esters and amides, for example, have been cleanly reduced to the corresponding aldehydes with the help of chlorotrimethylsilane in up to 64% yields[84]. Acyl halides and aliphatic amides have also been converted to aldehydes, albeit in modest yields[85].

5.0. Reduction of Carboxylic Acids and Carbamates to Alcohols

5.1. LiAlH_4 , NaBH_4 and BH_3

Lithium aluminum hydride (LiAlH_4), sodium borohydride (NaBH_4) and borane (BH_3) are common reducing agents used in organic transformations involving the conversion of carboxylic acids to the corresponding primary alcohols. Each of these three reducing agents operates by a different mechanism for their respective reductions. The choice of reducing agents for a particular transformation at hand is generally a topic covered in standard organic coursework. As such, we limit our discussion to just one example and illustrate the reduction of a carboxylic acid by BH_3 .



Scheme 11. Carboxylic acids reduction to alcohol with boranes.


The mechanism of reduction of carboxylic acids to alcohols with BH_3 occurs through the addition of a first borane equivalent across a $\text{C}=\text{O}$ bond ($53 \rightarrow 54$) followed by the coordination of a second borane equivalent to the newly-formed alcohol moiety ($54 \rightarrow 55$) and elimination of BH_2OBH_2 unit to lead to the formation of the primary alcohol product (56) (Scheme 11).

The electrochemical generation reducing agents that can accomplish transformations commonly performed by lithium aluminum hydride or borohydride species is not well-understood. Consequently, further electrochemical studies on the generation of such reducing agents will likely lead to new chemistries in the future.

5.3. Bouveault-Blanc Reduction[86, 87]

The Bouveault-Blanc reduction of aliphatic and aromatic esters is a method for the reduction of carboxylic acids to alcohols *via* a $1e^-$ reduction with a group IA metal such as sodium[87, 88].

The traditional Bouveault-Blanc reduction is the sequential Na reduction of a carboxylic acid derivative (**57**) in protic medium as described in Scheme 12. The utility of this chemical is rather limited as it occurs under extremely harsh reductive conditions with poor functional group tolerance and selectivity. The product of such a reaction is usually a mixture of alcoholic products. In a milder set of conditions, an ester can be reduced by SmI_2 in a polar aprotic solvent such as hexamethylphosphoramide (HMPA) instead of Na.



Scheme 12. Bouveault-Blanc Reduction Na reduction of carboxylate derivatives to primary alcohols.


5.4. Electrochemical Bouveault-Blanc Reduction

Since the mechanism of this transformation occurs via sequential $1e^-$ reductions, it was translatable to electrochemical conditions. Lam and coworkers have now adapted this chemistry to an electrode-driven process [89]. Study of reaction conditions have shown that the single-electron process occurs optimally in DMF and NMP, as THF and MeCN have reactivity issues due to the relatively narrow potential windows in the conditions. Temperature and electrode material effects have also been observed and yields of up to 50% were observed on Carbon electrodes at 130 °C in an electrolysis under reflux conditions.


5.5. Reduction of NHC Carboxylates with Silanes – Methanol Production [39] [40]

This particular chemistry was described in Section 3.3.1. The observed chemistry is limited by the strong Si-O bond formed[40].

5.6. Other Electrochemical Reductions [75]



Aromatic carboxylic acids (**60**) and ester derivatives (**62**) have been shown to be electrochemically reduced under cathodic conditions to the corresponding benzyl alcohols (**61** and **63** respectively) as depicted in Eq 9 and 10. Notably, the reduction also proceeds to break the aromaticity of benzoic acid in ethanol[75].




6.0. Reduction of Carboxylic Acids to Alkyls

6.1. Carboxylic Acid Reductase Tandem Catalysis

In section 4.1 we briefly touch on the reduction of carboxylic acids to aldehydes. In conjunction with an additional reducing enzyme, the intermediate aldehyde can be further reduced to an alkane[79, 80].

6.2. Barton McCombie Decarboxylation and Electrochemical Variant[90, 91]

The Barton-McCombie decarboxylation is the radical-induced loss of CO₂ from a thiohydroxamate ester (**67**) as shown in Scheme 13. A carboxylic acid can be derivatized to such an ester. In the presence of a radical initiator such as AIBN and a hydrogen donor such as trialkyl tin hydride, the formed carboxylate radical **68** decomposes with loss of CO₂.




Scheme 13. Mechanism for the Barton McCombie Decarboxylation reaction.


Carbonic and carbamic acids (Eq 13 and 14) are known to decarboxylate rather easily, whereas aliphatic acids do not (eq 15).

There are several synthetic applications of the acid decarboxylation, although a drawback of these conversions is the use of highly-toxic tin reagents. An electrochemical variant of this transformation was observed during the electrolysis of 1-isoquinoline carboxylic acid [92].

A series of related radical halodecarboxylation transformations are known.: the Hunsdiecker-Borodine [93-95] and the Simonini reaction[96]. These transformations convert carboxylate salts into the corresponding alkyl and aryl halides in the presence of oxidizing Br_2 and I_2 respectively. If X_2 ($\text{X}=\text{Cl}$, Br or I) is produced in an anodic process during an electrolysis, these reactions may lead to productive halodecarboxylation chemistries in the presence of carboxylic acid derivatives if migration of the anodic species to the cathode occurs.



6.3. Electrochemical[75]



This particular type of conversion is important for historical reasons, as it was the first example of electrochemical transformation ever reported. In 1834, Faraday reported oxidative decarboxylative coupling of acetic acid to ethane as a 2H^+ , 2e^- process[97]. Years later, Kolbe reported a generalized version of Faraday's experiments, known today as the Kolbe electrolysis[98].

As far as reductive processes are concerned, the Barton Decarboxylation remains state-of-the-art in the field, although benzylbenzoate (**70**) has been converted to benzoic acid (**71**) and toluene (**72**) (Eq 16). This particular reduction is likely a reduction of the alcohol portion of the ester and as such, it counts as a reduction of an alcohol to an alkyl rather than as a decarboxylation event.

Another example of reduction of a carboxylate species to the corresponding alkyls is the selective electrochemical conversion of CO_2 to methane as reported by Koval and coworkers [41]. In this work, the NHC-preactivation highlighted in Section 3.3.1 was utilized to gain access to an 8e^- reduced product: methane. The CO_2 is first converted to a zwitterionic carboxylate by reaction with a nucleophilic carbene. The CO_2 -adduct formed in this reaction was then exposed to reductive conditions in an electrochemical system in the presence of a Ni mediator in the presence of protons to effect the reductive chemistry.

7.0. Reduction of Aldehydes to Alcohols


7.1. NaBH_4 and LiAlH_4

We briefly described the reduction of carboxylic acids in Section 5.1. The reduction of aldehydes to alcohols is a repetition of the last step of the conversion of carboxylic acids to primary alcohols.

7.2. Crossed Cannizzaro Reaction: Formaldehyde as a Reducing Agent[99, 100]

The Cannizzaro Reaction is a transformation specific to aldehydes without an available $\alpha\text{-H}$ *ie* formaldehyde (**73**) or benzaldehyde and derivatives (**76**). In the presence of strong basic conditions such as NaOH (Eq 17 and 18), two equivalents of an aldehyde can undergo an intramolecular H transfer (Eq 17 TS) in which one aldehyde molecule is oxidized to a formate and another molecule is reduced to the corresponding alcohol. Taking advantage of the rate of this reaction, high yields of any alcohol-reduced products can be obtained by using an excess of formaldehyde, as the oxidation of formaldehyde to formic acid is a favorable process. This

process of using formaldehyde as a reducing agent is called the Crossed Cannizzaro reaction. A solvent-free variant of this reaction has been reported[101].



This particular transformation may become an important reaction in the field of solar fuels, as formaldehyde, a product on the CO_2 reduction pathway to methanol and methane, can *chemically* disproportionate under basic and thermal conditions to two different products, thus altering the product distributions. Thermal contributions to productive chemistry in electrochemical transformations for CO_2 reduction that can occur from resistive heating or illumination have been the subject of a recent report[102].

7.3. Silanes, [103] Boranes[104] and NHC Boranes[105]

In this section we enumerate some organocatalytic reduction chemistry with silanes and boranes as terminal reducing agents, as the transformations may become amenable to electrochemical conditions.

In a recent report, the hydrosilylation of carbon-heteroatom double bonds has been demonstrated in the presence of borohydride catalysts. Trace amounts of LiHBEt_3 and NaBH_4 have given high activity and good yields. In the case of this particular transformation, DFT calculations suggest the presence of a hypervalent Si intermediate[106, 107].

The activation of diphenylsilane by trace amounts of imidazolium salts (NHC precursors) has now been the protagonist of carbonyl reduction chemistry[103]. A similar hypervalent silicon active species has been invoked in the transformation, as the silicon intermediate can have strong Lewis acidic properties and also act as a strong hydride donor.

Pinacolborane has also been shown to catalytically reduce carbonyl compounds in the presence of NaOtBu as catalyst[104]. The active reducing agent in this reaction has been shown to be an *in-situ* generated trialkoxyborohydride.

Aldehyde reduction has also been demonstrated with 1,3-dimethylimidazol-2-ylidene trihydridoborane as the active hydride donor[105]. In the presence of silica gel, the conversion to

alcohols occurs smoothly and in good yields. Excitingly, this particular reactivity is specifically selective for aldehydes. Ketones are not reduced in the course of this chemical transformation.

7.4. Electrochemical Reduction: Reductive Dimerization[75]

The reductive electrochemistry of aldehydes is plagued with selectivity issues related to polymerization reactions[108, 109]. However, the key to long-sought productive C-C coupling reactions leading to the production of long-chain alkane fuels from CO₂ may lie in this particular reductive regime. Several examples of well-controlled polymerizations have been reported and we remain forward-looking to developments in the field[110, 111].

Of particular note in this direction is the work of Kanan *et al* [112] where the reduction of CO over nanocrystalline Cu yielded C₂ and C₃ products such as ethanol, acetate and *n*-propanol. A formyl radical coupling may be involved in the process.


7.5. Electrochemical Reduction on Cu Electrodes

Interestingly, a report of organic reduction of aldehydes over Cu shows great selectivity for alcohols without the formation of multimetric products[113]. The undivided electrolysis cell was connected to a 6V battery and was used to demonstrate the conversion of aliphatic and aromatic aldehydes to the corresponding alcohols.

8.0. Reduction of Aldehydes to Alkyls

8.1. Wolff-Kishner Reduction [114]

An aldehyde (**79**) reacts with hydrazine (**80**) to form a hydrazone (**81**). After the carbonyl activation step, under basic alcoholic conditions and heat, hydrazones can decompose to form the corresponding alkyls (**82**). The overall reaction sequence is known as the Wolff-Kishner reaction (Scheme 14). High temperatures are required, and the transformation is incompatible with hydrolysis-prone substrates such as esters, lactams, lactones, amides, etc. Examples of side-reactivity include the incomplete reduction of the carbonyl group to an alcohol rather than the alkane, as well as elimination reactions and double-bond isomerizations[114].




Scheme 14. Wolff-Kishner Reduction Mechanism.

8.2. Caglioti Reaction: Reduction of Tosylhydrazones[115]

A modification of the Wolff-Kishner Reduction, the Caglioti reaction occurs *via* the reduction of a tosylhydrazone[116]. Such reduction can be performed with mild borane reducing agents such as catecholboranes. Tosylhydrazide intermediates have been isolated[117] [118]. Other compatible reducing agents are lithium aluminum hydride [119] and the milder sodium cyanoborohydride[120].

8.3. Clemmensen Reduction[121]



The Clemmensen Reduction is the conversion of carbonyl compounds to the corresponding alkanes under harsh acidic conditions. The original report employed Zn/Hg amalgam, a $2e^-$ reducing agent. The aldehyde (83) undergoes two sequential $2e^-$ reduction steps through a dehydration step and two anionic intermediates **84** and **85**.



Scheme 15. Mechanism for the Clemmensen Reduction.

An interesting report highlights fuel production with Zn from formaldehyde *via* a Clemmensen reduction[122] [123].

8.5. Eschweiler-Clarke Reaction: Formaldehyde as an Oxidizing Agent[55]



Scheme 16. Mechanism for the Eschweiler-Clarke reaction.

The Eschweiler-Clark reaction is a reductive methylation reaction using an aldehyde (**88**) as an oxidizing agent. Eq 19 is an example of such a reaction with formaldehyde as a methylating agent for the corresponding amine (**87**).


The reaction proceeds *via* the formation of an iminium ion (**90**) from the corresponding aldehyde and amine starting materials. In the presence of formic acid as a reducing agent, the iminium is reduced to the corresponding methylamine. The reaction is an overall reductive amination.

This type of chemical transformation may come into play in attempts at using amines as pre-activating /bending agents for the capture of CO₂ as carbamates as described in Section 3.3.3. Upon the reduction of carbamates, both formaldehyde and formic acid could be possible reduction products under the electrochemical conditions. In the presence of heating from electrolytic cell resistance or illumination in the case of possible photochemical experiments, non-faradaic Eschweiler-Clark reactivity can lead to the formation of methylated intermediates.

The observed methanol products in the electrocatalytic reduction of CO₂ in the presence of mercaptopteridine[53] could be the product of chemical hydrolysis of an N-methyl intermediate under the electrochemical conditions. Independent isolation and electrochemical characterization of specific chemical intermediates could provide a rigorous reactivity assessment of chemical entities performing the observed transformations.

9.0. The Reduction of Alcohols to Alkyls

9.1. Barton-McCombie Deoxygenation[124]



The Barton McCombie Deoxygenation reaction is a radical-induced process that replaces the functional group of an alcohol (**90**) with hydrogen (Eq 20) to form an alkane (**92**). As such, it employs a H donor, usually a SnH, akin to the terminal reducing agent used in the decarboxylation reaction carrying the same name (Section 6.2). The first step in this transformation is the formation of a thioester derivative of the starting material, which can then undergo reduction with the tin reducing agent. The downside of this chemistry is the use of highly toxic tin reagents.

In recent reports, a photoredox process has been reported[125]. In a similar vein, the Cu-catalyzed reduction of alkyl triflates to the corresponding alkane, has also been reported[126]. Applications in total synthesis of this reaction have been reviewed by Faghihi and coworkers[127].

Several other deoxygenation methods have been reported and have been reviewed elsewhere[128, 129]. For the purpose of the current discussion we describe one electrochemical instance of such transformation in the following section.

9.2. Electrochemical methods

Shono has long reported the cathodic deoxygenation of mesylates as an alternative to liquid ammonia reductions of mesylates and triflates[128, 130]. The electrochemical method was attractive in that it had a good functional group tolerance: tolerating carboxylic acids, nitriles, olefins and epoxide functionality. Unfortunately electrolytic methods have not yet been adopted as a mainstream tool of organic synthesis.

10. Conclusion

Although a long-standing interest for many decades, the field of CO₂ reduction has not yet advanced to the stage of technological implementation. From an organic chemistry perspective, the solar fuels field will likely contribute a wealth of knowledge regarding the electrochemical generation of selective and functional-group tolerant reducing agents. As such, efforts in the electrochemical conversions of CO₂ to solar fuels may provide mild conditions for transformations related to carboxylic acids and intermediates.

CO₂ capture and subsequent reduction chemistry has unfortunately not yet been fully elucidated in terms of electrochemical regimes and we remain mindful of coupled chemical processes related to the chemical reactivity of possible intermediates. We look forward to the concept of chemical preactivation of CO₂ as an energy-relevant molecule as we propose that these transformations could be approached as an exercise in the electrochemical reduction of carboxylates and related further-reduced species.

Acknowledgements

This material is based upon work performed by the Joint Center for Artificial Photosynthesis, a DOE Energy Innovation Hub, supported through the Office of Science of the U.S. Department of Energy under Award Number DE-SC0004993. The authors would like to thank Mr. Michael Lichterman, Ms. Maeghan Deegan, Dr. David K. Romney, Dr. David Lacy, Ms. Shannon Carpenter, Professors Carl Koval, Jonas Peters, T. Don Tilley, Clifford Kubiak and Nathan Lewis for useful discussions.

- [1] A. Q. Fenwick, J. M. Gregore, and O. R. Luca, "Electrocatalytic Reduction of Nitrogen and Carbon Dioxide to Chemical Fuels: Challenges and Opportunities for a Solar Fuel Device," *Journal of Photochemistry and Photobiology B: Biology*, 2014, 10.1016/j.jphotobiol.2014.12.019.

- [2] D. W. C. MacMillan, "The advent and development of organocatalysis," *Nature*, vol. 455, pp. 304-308, 2008.
- [3] R. Francke and R. D. Little, "Redox catalysis in organic electrosynthesis: basic principles and recent developments," *Chemical Society Reviews*, vol. 43, pp. 2492-2521, 2014.
- [4] J. B. Broderick, "Coenzymes and Cofactors," *eLS*, 2000.
- [5] Y. Oh and X. Hu, "Organic molecules as mediators and catalysts for photocatalytic and electrocatalytic CO₂ reduction," *Chemical Society Reviews*, vol. 42, pp. 2253-2261, 2013.
- [6] J. O. M. Bockris and J. C. Wass, "The Photoelectrocatalytic Reduction of Carbon Dioxide," *Journal of The Electrochemical Society*, vol. 136, pp. 2521-2528, September 1, 1989 1989.
- [7] B. A. Rosen, A. Salehi-Khojin, M. R. Thorson, W. Zhu, D. T. Whipple, P. J. A. Kenis, *et al.*, "Ionic Liquid-Mediated Selective Conversion of CO₂ to CO at Low Overpotentials," *Science*, vol. 334, pp. 643-644, November 4, 2011 2011.
- [8] E. Barton Cole, P. S. Lakkaraju, D. M. Rampulla, A. J. Morris, E. Abelev, and A. B. Bocarsly, "Using a One-Electron Shuttle for the Multielectron Reduction of CO₂ to Methanol: Kinetic, Mechanistic, and Structural Insights," *Journal of the American Chemical Society*, vol. 132, pp. 11539-11551, 2010/08/25 2010.
- [9] S. G. Jadhav, P. D. Vaidya, B. M. Bhanage, and J. B. Joshi, "Catalytic carbon dioxide hydrogenation to methanol: A review of recent studies," *Chemical Engineering Research and Design*, 2014.
- [10] J. E. Lyon, L. E. Rennick, and J. L. Burmeister, "Use of Transition Metal Complexes as Selective Hydrogenation Catalysts," *Product R&D*, vol. 9, pp. 2-20, 1970/03/01 1970.
- [11] R. Crabtree, "Iridium compounds in catalysis," *Accounts of Chemical Research*, vol. 12, pp. 331-337, 1979.
- [12] Y. Wei, X. Wu, C. Wang, and J. Xiao, "Transfer hydrogenation in aqueous media," *Catalysis Today*, 2014.
- [13] J. Melville, C. Liu, E. Wang, G. Zhou, and R. Triano, "Chemoselectivity in Transfer-Hydrogenation Reactions," 2014.
- [14] O. R. Luca, T. Wang, S. J. Konezny, V. S. Batista, and R. H. Crabtree, "DDQ as an electrocatalyst for amine dehydrogenation, a model system for virtual hydrogen storage," *New J. Chem.*, vol. 35, pp. 998-999, 2011.
- [15] O. R. Luca, D. L. Huang, M. K. Takase, and R. H. Crabtree, "Redox-active cyclopentadienyl Ni complexes with quinoid N-heterocyclic carbene ligands for the electrocatalytic hydrogen release from chemical fuels," *New J. Chem.*, vol. 37, pp. 3402-3405, 2013.
- [16] O. R. Luca, B. A. Thompson, M. K. Takase, and R. H. Crabtree, "Synthesis and electrochemistry of a series of cyclopentadienyl Ni N-heterocyclic carbene compounds," *Journal of Organometallic Chemistry*, vol. 730, pp. 79-83, 2013.
- [17] B. Kumar, M. Llorente, J. Froehlich, T. Dang, A. Sathrum, and C. P. Kubiak, "Photochemical and photoelectrochemical reduction of CO₂," *Annual review of physical chemistry*, vol. 63, pp. 541-569, 2012.
- [18] A. A. Peterson, F. Abild-Pedersen, F. Studt, J. Rossmeisl, and J. K. Nørskov, "How copper catalyzes the electroreduction of carbon dioxide into hydrocarbon fuels," *Energy & Environmental Science*, vol. 3, pp. 1311-1315, 2010.
- [19] H. A. Hansen, J. B. Varley, A. A. Peterson, and J. K. Nørskov, "Understanding Trends in the Electrocatalytic Activity of Metals and Enzymes for CO₂ Reduction to CO," *The Journal of Physical Chemistry Letters*, vol. 4, pp. 388-392, 2013/02/07 2013.
- [20] A. Dhingra, A. R. Portis, and H. Daniell, "Enhanced translation of a chloroplast-expressed RbcS gene restores small subunit levels and photosynthesis in nuclear RbcS antisense plants,"

- Proceedings of the National Academy of Sciences of the United States of America*, vol. 101, pp. 6315-6320, April 20, 2004 2004.
- [21] U. Feller, I. Anders, and T. Mae, "Rubiscolytics: fate of Rubisco after its enzymatic function in a cell is terminated," *Journal of Experimental Botany*, vol. 59, pp. 1615-1624, May 1, 2008 2008.
- [22] B. E. Maden, "Tetrahydrofolate and tetrahydromethanopterin compared: functionally distinct carriers in C1 metabolism," *Biochemical Journal*, vol. 350, pp. 609-629, 2000.
- [23] P. Weimer and J. Zeikus, "One carbon metabolism in methanogenic bacteria," *Archives of microbiology*, vol. 119, pp. 49-57, 1978.
- [24] L. G. Ujungdahl, "Comparative Biochemistry of C1-Carriers," in *Microbial Growth on C1 Compounds*, ed: Springer, 1987, pp. 121-128.
- [25] R. K. Thauer, R. Hedderich, and R. Fischer, "Reactions and enzymes involved in methanogenesis from CO₂ and H₂," in *Methanogenesis*, ed: Springer, 1993, pp. 209-252.
- [26] V. Müller, M. Blaut, and G. Gottschalk, "Bioenergetics of methanogenesis," in *Methanogenesis*, ed: Springer, 1993, pp. 360-406.
- [27] P. Raemakers-Franken, A. Kortstee, C. Van der Drift, and G. Vogels, "Methanogenesis involving a novel carrier of C1 compounds in Methanogenium tationis," *Journal of bacteriology*, vol. 172, pp. 1157-1159, 1990.
- [28] J. G. Ferry, "Methane from acetate," *Journal of bacteriology*, vol. 174, pp. 5489-5495, 1992.
- [29] F. Glorius and S. Bellemin-Laponnaz, *N-heterocyclic carbenes in transition metal catalysis*: Springer, 2007.
- [30] S. P. Nolan, *N-Heterocyclic carbenes in synthesis*: John Wiley & Sons, 2006.
- [31] W. A. Herrmann and C. Koecher, "N-Heterocyclic Carbenes," *Angewandte Chemie International Edition in English*, vol. 36, pp. 2162-2187, 1997.
- [32] A. M. Voutchkova, M. Feliz, E. Clot, O. Eisenstein, and R. H. Crabtree, "Imidazolium carboxylates as versatile and selective N-heterocyclic carbene transfer agents: Synthesis, mechanism, and applications," *Journal of the American Chemical Society*, vol. 129, pp. 12834-12846, 2007.
- [33] B. R. Van Ausdall, J. L. Glass, K. M. Wiggins, A. M. Aarif, and J. Louie, "A systematic investigation of factors influencing the decarboxylation of imidazolium carboxylates," *The Journal of organic chemistry*, vol. 74, pp. 7935-7942, 2009.
- [34] B. R. Van Ausdall, N. F. Poth, V. A. Kincaid, A. M. Arif, and J. Louie, "Imidazolidene Carboxylate Bound MBPh₄ Complexes (M= Li, Na) and Their Relevance in Transcarboxylation Reactions," *The Journal of organic chemistry*, vol. 76, pp. 8413-8420, 2011.
- [35] B. R. Van Ausdall, "An investigation of N-heterocyclic carbene carboxylates: Insight into decarboxylation, a transcarboxylation reaction, and synthesis of hydrogen bonding precursors," The University of Utah, 2012.
- [36] H. Zhou, W.-Z. Zhang, C.-H. Liu, J.-P. Qu, and X.-B. Lu, "CO₂ adducts of N-heterocyclic carbenes: thermal stability and catalytic activity toward the coupling of CO₂ with epoxides," *The Journal of organic chemistry*, vol. 73, pp. 8039-8044, 2008.
- [37] E. E. Finney, K. A. Ogawa, and A. J. Boydston, "Organocatalyzed Anodic Oxidation of Aldehydes," *Journal of the American Chemical Society*, vol. 134, pp. 12374-12377, 2012/08/01 2012.
- [38] S. N. Riduan, Y. Zhang, and J. Y. Ying, "Conversion of Carbon Dioxide into Methanol with Silanes over N-Heterocyclic Carbene Catalysts," *Angewandte Chemie*, vol. 121, pp. 3372-3375, 2009.
- [39] F. Huang, G. Lu, L. Zhao, H. Li, and Z.-X. Wang, "The Catalytic Role of N-Heterocyclic Carbene in a Metal-Free Conversion of Carbon Dioxide into Methanol: A Computational Mechanism Study," *Journal of the American Chemical Society*, vol. 132, pp. 12388-12396, 2010/09/08 2010.
- [40] S. N. Riduan and Y. Zhang, "Recent developments in carbon dioxide utilization under mild conditions," *Dalton Transactions*, vol. 39, pp. 3347-3357, 2010.

- [41] O. R. Luca, C.C.L. McCrory, Dalleska N.F., Koval C.A. "The Selective Electrochemical Conversion of Carbon Dioxide to Methane," *Journal of the Electrochemical Society*, DOI: 10.1149/2.0371507jes 2015.
- [42] S. M. Ibrahim Al-Rafia, A. C. Malcolm, S. K. Liew, M. J. Ferguson, R. McDonald, and E. Rivard, "Intercepting low oxidation state main group hydrides with a nucleophilic N-heterocyclic olefin," *Chemical Communications*, vol. 47, pp. 6987-6989, 2011.
- [43] Y. Wang, M. Y. Abraham, R. J. Gilliard, D. R. Sexton, P. Wei, and G. H. Robinson, "N-Heterocyclic Olefin Stabilized Borenium Cations," *Organometallics*, vol. 32, pp. 6639-6642, 2013/11/25 2013.
- [44] Y.-B. Jia, Y.-B. Wang, W.-M. Ren, T. Xu, J. Wang, and X.-B. Lu, "Mechanistic Aspects of Initiation and Deactivation in N-Heterocyclic Olefin Mediated Polymerization of Acrylates with Alane as Activator," *Macromolecules*, vol. 47, pp. 1966-1972, 2014/03/25 2014.
- [45] Y.-B. Wang, Y.-M. Wang, W.-Z. Zhang, and X.-B. Lu, "Fast CO₂ Sequestration, Activation, and Catalytic Transformation Using N-Heterocyclic Olefins," *Journal of the American Chemical Society*, vol. 135, pp. 11996-12003, 2013/08/14 2013.
- [46] P. Danckwerts, "The reaction of CO₂ with ethanolamines," *Chemical Engineering Science*, vol. 34, pp. 443-446, 1979.
- [47] G. Sartori and D. W. Savage, "Sterically hindered amines for carbon dioxide removal from gases," *Industrial & Engineering Chemistry Fundamentals*, vol. 22, pp. 239-249, 1983.
- [48] P. D. Vaidya and E. Y. Kenig, "CO₂-Alkanolamine Reaction Kinetics: A Review of Recent Studies," *Chemical Engineering & Technology*, vol. 30, pp. 1467-1474, 2007.
- [49] A. Tlili, E. Blondiaux, X. Frogneux, and T. Cantat, "Reductive functionalization of CO₂ with amines: an entry to formamide, formamidine and methylamine derivatives," *Green Chemistry*, vol. 17, pp. 157-168, 2015.
- [50] C. Wu, H. Cheng, R. Liu, Q. Wang, Y. Hao, Y. Yu, *et al.*, "Synthesis of urea derivatives from amines and CO₂ in the absence of catalyst and solvent," *Green Chemistry*, vol. 12, pp. 1811-1816, 2010.
- [51] M. Z. Kamrath, R. A. Relph, and M. A. Johnson, "Vibrational predissociation spectrum of the carbamate radical anion, C₅H₅N-CO₂⁻, generated by reaction of pyridine with (CO₂)⁻," *Journal of the American Chemical Society*, vol. 132, pp. 15508-15511, 2010.
- [52] A. J. Morris, R. T. McGibbon, and A. B. Bocarsly, "Electrocatalytic Carbon Dioxide Activation: The Rate-Determining Step of Pyridinium-Catalyzed CO₂ Reduction," *ChemSusChem*, vol. 4, pp. 191-196, 2011.
- [53] D. Xiang, D. Magana, and R. B. Dyer, "CO₂ Reduction Catalyzed by Mercaptopteridine on Glassy Carbon," *Journal of the American Chemical Society*, vol. 136, pp. 14007-14010, 2014/10/08 2014.
- [54] H.-C. Hsu and D.-R. Hou, "Reduction of 1-pyrrolyl and 1-indolyl carbamates to hemiaminals," *Tetrahedron Letters*, vol. 50, pp. 7169-7171, 12/23/ 2009.
- [55] J. J. Li, "Eschweiler-Clarke reductive alkylation of amines," in *Name Reactions*, ed: Springer, 2014, pp. 235-236.
- [56] Q.-Y. Bi, J.-D. Lin, Y.-M. Liu, X.-L. Du, J.-Q. Wang, H.-Y. He, *et al.*, "An Aqueous Rechargeable Formate-Based Hydrogen Battery Driven by Heterogeneous Pd Catalysis," *Angewandte Chemie International Edition*, vol. 53, pp. 13583-13587, 2014.
- [57] O. R. Luca, "Virtual Hydrogen Storage Strategies for the Production and Storage of Hydrogen," YALE UNIVERSITY, 2013.
- [58] R. H. Crabtree, "Hydrogen storage in liquid organic heterocycles," *Energy & Environmental Science*, vol. 1, pp. 134-138, 2008.
- [59] C.-H. Lim, A. M. Holder, J. T. Hynes, and C. B. Musgrave, "Roles of the Lewis Acid and Base in the Chemical Reduction of CO₂ Catalyzed by Frustrated Lewis Pairs," *Inorganic Chemistry*, vol. 52, pp. 10062-10066, 2013/09/03 2013.

- [60] A. Berkefeld, W. E. Piers, and M. Parvez, "Tandem frustrated Lewis pair/tris (pentafluorophenyl) borane-catalyzed deoxygenative hydrosilylation of carbon dioxide," *Journal of the American Chemical Society*, vol. 132, pp. 10660-10661, 2010.
- [61] D. W. Stephan, "Frustrated Lewis Pairs: From Concept to Catalysis," *Accounts of Chemical Research*, 2014.
- [62] A. Berkefeld, W. E. Piers, and M. Parvez, "Tandem Frustrated Lewis Pair/Tris(pentafluorophenyl)borane-Catalyzed Deoxygenative Hydrosilylation of Carbon Dioxide," *Journal of the American Chemical Society*, vol. 132, pp. 10660-10661, 2010/08/11 2010.
- [63] M. Khandelwal and R. J. Wehmschulte, "Deoxygenative Reduction of Carbon Dioxide to Methane, Toluene, and Diphenylmethane with $[\text{Et}_2\text{Al}]^+$ as Catalyst," *Angewandte Chemie International Edition*, vol. 51, pp. 7323-7326, 2012.
- [64] L. J. Hounjet and D. W. Stephan, "Hydrogenation by Frustrated Lewis Pairs: Main Group Alternatives to Transition Metal Catalysts?," *Organic Process Research & Development*, vol. 18, pp. 385-391, 2014/03/21 2014.
- [65] T. Mahdi and D. W. Stephan, "Enabling Catalytic Ketone Hydrogenation by Frustrated Lewis Pairs," *Journal of the American Chemical Society*, vol. 136, pp. 15809-15812, 2014/11/12 2014.
- [66] A. Y. Houghton, J. Hurmalainen, A. Mansikkamäki, W. E. Piers, and H. M. Tuononen, "Direct observation of a borane-silane complex involved in frustrated Lewis-pair-mediated hydrosilylations," *Nat Chem*, vol. 6, pp. 983-988, 11//print 2014.
- [67] X. Feng and H. Du, "Metal-free asymmetric hydrogenation and hydrosilylation catalyzed by frustrated Lewis pairs," *Tetrahedron Letters*, vol. 55, pp. 6959-6964, 12/17/ 2014.
- [68] D. H. Gibson, "The Organometallic Chemistry of Carbon Dioxide," *Chemical Reviews*, vol. 96, pp. 2063-2096, 1996/01/01 1996.
- [69] W. Leitner, "The coordination chemistry of carbon dioxide and its relevance for catalysis: a critical survey," *Coordination chemistry reviews*, vol. 153, pp. 257-284, 1996.
- [70] E. E. Benson, C. P. Kubiak, A. J. Sathrum, and J. M. Smieja, "Electrocatalytic and homogeneous approaches to conversion of CO_2 to liquid fuels," *Chemical Society Reviews*, vol. 38, pp. 89-99, 2009.
- [71] O. R. Luca and R. H. Crabtree, "Redox-active ligands in catalysis," *Chemical Society Reviews*, vol. 42, pp. 1440-1459, 2013.
- [72] E. Balaraman, C. Gunanathan, J. Zhang, L. J. W. Shimon, and D. Milstein, "Efficient hydrogenation of organic carbonates, carbamates and formates indicates alternative routes to methanol based on CO_2 and CO ," *Nat Chem*, vol. 3, pp. 609-614, 08//print 2011.
- [73] J. M. Smieja, M. D. Sampson, K. A. Grice, E. E. Benson, J. D. Froehlich, and C. P. Kubiak, "Manganese as a Substitute for Rhenium in CO_2 Reduction Catalysts: The Importance of Acids," *Inorganic Chemistry*, vol. 52, pp. 2484-2491, 2013/03/04 2013.
- [74] M. D. Sampson, A. D. Nguyen, K. A. Grice, C. E. Moore, A. L. Rheingold, and C. P. Kubiak, "Manganese Catalysts with Bulky Bipyridine Ligands for the Electrocatalytic Reduction of Carbon Dioxide: Eliminating Dimerization and Altering Catalysis," *Journal of the American Chemical Society*, vol. 136, pp. 5460-5471, 2014/04/09 2014.
- [75] S. Torii, *Electroorganic reduction synthesis* vol. 1: Kodansha Tokyo, 2006.
- [76] G. Silvestri, S. Gambino, and G. Filardo, "Electrochemical Syntheses Involving Carbon Dioxide," in *Enzymatic and Model Carboxylation and Reduction Reactions for Carbon Dioxide Utilization*, ed: Springer, 1990, pp. 101-127.
- [77] J. Song, E. L. Klein, F. Neese, and S. Ye, "The Mechanism of Homogeneous CO_2 Reduction by Ni (cyclam): Product Selectivity, Concerted Proton-Electron Transfer and C-O Bond Cleavage," *Inorganic chemistry*, vol. 53, pp. 7500-7507, 2014.

- [78] J. D. Froehlich and C. P. Kubiak, "Homogeneous CO₂ Reduction by Ni (cyclam) at a Glassy Carbon Electrode," *Inorganic chemistry*, vol. 51, pp. 3932-3934, 2012.
- [79] M. K. Akhtar, N. J. Turner, and P. R. Jones, "Carboxylic acid reductase is a versatile enzyme for the conversion of fatty acids into fuels and chemical commodities," *Proceedings of the National Academy of Sciences*, vol. 110, pp. 87-92, 2013.
- [80] H. WHITE, G. STROBL, R. FEICHT, and H. SIMON, "Carboxylic acid reductase: a new tungsten enzyme catalyses the reduction of non-activated carboxylic acids to aldehydes," *European Journal of Biochemistry*, vol. 184, pp. 89-96, 1989.
- [81] C. Das Neves Gomes, O. Jacquet, C. Villiers, P. Thuéry, M. Ephritikhine, and T. Cantat, "A Diagonal Approach to Chemical Recycling of Carbon Dioxide: Organocatalytic Transformation for the Reductive Functionalization of CO₂," *Angewandte Chemie International Edition*, vol. 51, pp. 187-190, 2012.
- [82] L. Schmid, A. Canonica, and A. Baiker, "Ruthenium-catalysed formylation of amines with dense carbon dioxide as C1-source," *Applied catalysis A: general*, vol. 255, pp. 23-33, 2003.
- [83] L. Schmid, M. S. Schneider, D. Engel, and A. Baiker, "Formylation with "supercritical" CO₂: efficient ruthenium-catalyzed synthesis of N-formylmorpholine," *Catalysis letters*, vol. 88, pp. 105-113, 2003.
- [84] P. R. Goetz-Schatowitz, G. Struth, J. Voss, and G. Wiegand, "Elektroreduktion organischer Verbindungen. 22. Darstellung aromatischer Aldehyde durch Elektroreduktion von Estern und Amiden in Gegenwart von Chlortrimethylsilan," *Journal für Praktische Chemie/Chemiker-Zeitung*, vol. 335, pp. 230-234, 1993.
- [85] G. A. Urove, D. G. Peters, and M. S. Mubarak, "Production of aldehydes via electrochemical reduction of acyl halides at mercury and carbon cathodes in acetonitrile," *The Journal of Organic Chemistry*, vol. 57, pp. 786-790, 1992.
- [86] B.-I. Seo, L. K. Wall, H. Lee, J. W. Buttrum, and D. E. Lewis, "An Improved Practical Synthesis of Isomerically Pure 3-endo-(p-Methoxybenzyl) isoborneol," *Synthetic communications*, vol. 23, pp. 15-22, 1993.
- [87] B. S. Bodnar and P. F. Vogt, "An Improved Bouveault–Blanc Ester Reduction with Stabilized Alkali Metals," *The Journal of organic chemistry*, vol. 74, pp. 2598-2600, 2009.
- [88] J. J. Li, *Name reactions: A collection of detailed mechanisms and synthetic applications*: Springer, 2010.
- [89] K. Lam and I. E. Markó, "Toluates: unexpectedly versatile reagents," *Tetrahedron*, vol. 65, pp. 10930-10940, 12/26/ 2009.
- [90] D. H. Barton, D. Crich, and W. B. Motherwell, "New and improved methods for the radical decarboxylation of acids," *J. Chem. Soc., Chem. Commun.*, pp. 939-941, 1983.
- [91] M. S. Baird, D. Bellus, N. Chessum, S. Couty, A. Dzielendziak, P. Eilbracht, *et al.*, *Science of Synthesis: Houben-Weyl Methods of Molecular Transformations Vol. 48: Alkanes*: Georg Thieme Verlag, 2014.
- [92] C. M. Sánchez-Sánchez, E. Expósito, B. Batanero, V. Montiel, F. Barba, and A. Aldaz, "Electrochemical reduction of the anion of 1-isoquinolinecarboxylic acid: an unexpected reaction of cathodic decarboxylation," *Electrochemistry communications*, vol. 6, pp. 595-599, 2004.
- [93] A. Borodine, "Ueber Bromvaleriansäure und Brombuttersäure," *Justus Liebigs Annalen der Chemie*, vol. 119, pp. 121-123, 1861.
- [94] H. Hunsdiecker and C. Hunsdiecker, "Über den Abbau der Salze aliphatischer Säuren durch Brom," *Berichte der deutschen chemischen Gesellschaft (A and B Series)*, vol. 75, pp. 291-297, 1942.

- [95] R. G. Johnson and R. K. Ingham, "The degradation of carboxylic acid salts by means of halogen-The Hunsdiecker reaction," *Chemical Reviews*, vol. 56, pp. 219-269, 1956.
- [96] A. Simonini, "Über den Abbau der fetten Säuren zu kohlenstoffärmeren Alkoholen," *Monatshefte für Chemie/Chemical Monthly*, vol. 13, pp. 320-325, 1892.
- [97] M. Faraday, "Experimental Researches in Electricity. Seventh Series.," *Phil. Trans. R. Soc. London*, vol. 124, pp. 77-122, 1834.
- [98] H. Kolbe, "XVII.—Researches on the electrolysis of organic compounds," *QJ Chem. Soc.*, vol. 2, pp. 157-184, 1850.
- [99] D. Davidson and M. T. Bogert, "The preparation of aromatic alcohols by the crossed Cannizzaro reaction with formaldehyde," *Journal of the American Chemical Society*, vol. 57, pp. 905-905, 1935.
- [100] J. J. Li, "Cannizzaro reaction," in *Name Reactions*, ed: Springer, 2014, pp. 106-107.
- [101] M. M. Mojtahedi, E. Akbarzadeh, R. Sharifi, and M. S. Abaee, "Lithium Bromide as a Flexible, Mild, and Recyclable Reagent for Solvent-Free Cannizzaro, Tishchenko, and Meerwein-Ponndorf-Verley Reactions," *Organic Letters*, vol. 9, pp. 2791-2793, 2007/07/01 2007.
- [102] A. Q. Fenwick and O. R. Luca, "Thermal Decomposition of Formic Acid under Electrochemical Conditions for CO₂ Reduction as a Source of CO " *J Photochem Photobiol B*, vol. submitted.
- [103] Q. Zhao, D. P. Curran, M. Malacria, L. Fensterbank, J.-P. Goddard, and E. Lacôte, "NHC-Catalyzed Chemo- and Regioselective Hydrosilylation of Carbonyl Derivatives," *Synlett*, vol. 2012, pp. 433-437.
- [104] I. P. Query, P. A. Squier, E. M. Larson, N. A. Isley, and T. B. Clark, "Alkoxide-Catalyzed Reduction of Ketones with Pinacolborane," *The Journal of Organic Chemistry*, vol. 76, pp. 6452-6456, 2011/08/05 2011.
- [105] T. Taniguchi and D. P. Curran, "Silica Gel Promotes Reductions of Aldehydes and Ketones by N-Heterocyclic Carbene Boranes," *Organic Letters*, vol. 14, pp. 4540-4543, 2012/09/07 2012.
- [106] M. G. Manas, L. S. Sharninghausen, D. Balcells, and R. H. Crabtree, "Experimental and computational studies of borohydride catalyzed hydrosilylation of a variety of C = O and C = N functionalities including esters, amides and heteroarenes," *New Journal of Chemistry*, vol. 38, pp. 1694-1700, 2014.
- [107] R. Corriu, G. Lanneau, and M. Perrot, "Facile reduction of acid chlorides into aldehydes using hypervalent silicon hydrides," *Tetrahedron letters*, vol. 29, pp. 1271-1274, 1988.
- [108] A. W. Cooke and K. Wagener, "An investigation of polymerization via reductive coupling of carbonyls," *Macromolecules*, vol. 24, pp. 1404-1407, 1991.
- [109] A. S. Amarasekara, B. Wiredu, and A. Razzaq, "Vanillin based polymers: I. An electrochemical route to polyvanillin," *Green Chemistry*, vol. 14, pp. 2395-2397, 2012.
- [110] W. R. Fawcett and A. Lasia, "The electroreduction of aromatic aldehydes in aprotic solvents," *Canadian Journal of Chemistry*, vol. 59, pp. 3256-3260, 1981.
- [111] T. Hirao, "Catalytic Reductive Coupling of Carbonyl Compounds—The Pinacol Coupling Reaction and Beyond," in *Metal Catalyzed Reductive C—C Bond Formation*, ed: Springer, 2007, pp. 53-75.
- [112] C. W. Li, J. Ciston, and M. W. Kanan, "Electroreduction of carbon monoxide to liquid fuel on oxide-derived nanocrystalline copper," *Nature*, vol. 508, pp. 504-507, 2014.
- [113] S. Velázquez-Olvera, H. Salgado-Zamora, M.-E. Campos-Aldrete, A. Reyes-Arellano, C. Pérez-González, and M. Velázquez-Ponce, "A convenient electrolytic process for the reduction of aldehydes," *Green Chemistry Letters and Reviews*, vol. 7, pp. 296-300, 2014.

- [114] S. G. Newman, L. Gu, C. Lesniak, G. Victor, F. Meschke, L. Abahmane, *et al.*, "Rapid Wolff–Kishner reductions in a silicon carbide microreactor," *Green Chemistry*, vol. 16, pp. 176-180, 2014.
- [115] V. K. Aggarwal, D. Bellus, S. J. Collier, M. Dabdoub, W. Eberbach, L. Fisera, *et al.*, *Science of Synthesis: Houben-Weyl Methods of Molecular Transformations Vol. 27: Heteroatom Analogues of Aldehydes and Ketones*: Georg Thieme Verlag, 2014.
- [116] L. Caglioti, "The reduction of tosylhydrazones and of acyl tosylhydrazides," *Tetrahedron*, vol. 22, pp. 487-493, 1966.
- [117] G. W. Kabalka, D. T. Yang, and J. D. Baker, "Deoxygenation of α,β -unsaturated aldehydes and ketones via the catecholborane reduction of the corresponding tosylhydrazones," *The Journal of Organic Chemistry*, vol. 41, pp. 574-575, 1976.
- [118] J. M. Khurana, B. Kandpal, P. Sharma, and M. Gupta, "A novel method of reduction of $C=N$ -group in hydrazones, phenylhydrazones, azines, and tosylhydrazones by Mg–methanol," *Monatshefte für Chemie-Chemical Monthly*, pp. 1-4, 2014.
- [119] L. Caglioti and M. Magi, "The reaction of tosylhydrazones with lithium aluminium hydride," *Tetrahedron*, vol. 19, pp. 1127-1131, 1963.
- [120] V. P. Miller, D. Y. Yang, T. M. Weigel, O. Han, and H. W. Liu, "Studies of the mechanistic diversity of sodium cyanoborohydride reduction of tosylhydrazones," *The Journal of Organic Chemistry*, vol. 54, pp. 4175-4188, 1989.
- [121] J. J. Li, "Clemmensen reduction," in *Name Reactions*, ed: Springer, 2014, pp. 153-154.
- [122] J. Myllyoja, J. Jakkula, P. Aalto, E. Koivusalmi, and J. Moilanen, "Process for producing a hydrocarbon component," ed: Google Patents, 2014.
- [123] J. M. Stafford, "Producing Fuel from Carbon Dioxide Using Zinc," *2014 NCUR*, 2014.
- [124] J. J. Li, "Barton–McCombie deoxygenation," in *Name Reactions*, ed: Springer, 2014, pp. 28-29.
- [125] D. Rackl, V. Kais, P. Kreitmeier, and O. Reiser, "Visible light photoredox-catalyzed deoxygenation of alcohols," *Beilstein journal of organic chemistry*, vol. 10, pp. 2157-2165, 2014.
- [126] H. Dang, N. Cox, and G. Lalic, "Copper-Catalyzed Reduction of Alkyl Triflates and Iodides: An Efficient Method for the Deoxygenation of Primary and Secondary Alcohols," *Angewandte Chemie International Edition*, vol. 53, pp. 752-756, 2014.
- [127] M. M. Heravi, A. Bakhtiari, and Z. Faghihi, "Applications of Barton-McCombie reaction in total syntheses," *Current Organic Synthesis*, vol. 11, pp. 787-823, 2014.
- [128] W. Hartwig, "Modern methods for the radical deoxygenation of alcohols," *Tetrahedron*, vol. 39, pp. 2609-2645, 1983.
- [129] H. Alper and M. Sališová, "The iron carbonyl induced deoxygenation of alcohols," *Tetrahedron Letters*, vol. 21, pp. 801-804, 1980.
- [130] T. Shono, Y. Matsumura, K. Tsubata, and S. Yoshihiro, "Selective transformation of aliphatic alcohols to alkanes by electroreductive method," *Tetrahedron Letters*, vol. 20, pp. 2157-2160, 1979.

We survey the organic chemistry involved in the reduction of CO₂ and related intermediates to chemical fuels.

We describe chemical reactions of CO₂ with organic molecules to yield carboxylic acid derivatives that can get further reduced.

We point out reactions such as the Cross Canizzaro and the Eschweiler Clark as possible non-faradaic chemical reactions contributing to observed product distributions in the production of solar fuels from CO₂.

